THE WEATHER Of ONTARIO AND QUEBEC



GRAPHIC AREA FORECAST 33



THE WEATHER OF ONTARIO AND QUEBEC

GRAPHIC AREA FORECAST 33

by Ross Klock Gilles Simard John Mullock



Copyright

Copyright © 2002 NAV CANADA. All rights reserved. No part of this document may be reproduced in any form, including photocopying or transmission electronically to any computer, without prior written consent of NAV CANADA. The information contained in this document is confidential and proprietary to NAV CANADA and may not be used or disclosed except as expressly authorized in writing by NAV CANADA.

Trademarks

Product names mentioned in this document may be trademarks or registered trademarks of their respective companies and are hereby acknowledged.

Relief Maps

Copyright © 2000. Government of Canada with permission from Natural Resources Canada



Design and illustration by Ideas in Motion Kelowna, British Columbia ph: (250) 717-5937 ideasinmotion@shaw.ca

The Weather of Ontario-Quebec Graphic Area Forecast 33 Ontario-Quebec

Preface

For NAV CANADA's Flight Service Specialists (FSS), providing weather briefings to help pilots navigate through the day to day fluctuations in the weather is a critical role. While available weather products are becoming increasingly more sophisticated and at the same time more easily understood, an understanding of local and regional climatological patterns is essential to the effective performance of this role.

This Ontario-Quebec Local Area Knowledge Aviation Weather manual is one of a series of six publications prepared by Meteorological Services of Canada for NAV CANADA. Each of the six manuals corresponds to a specific graphic forecast area (GFA) Domain, with the exception of the Nunavut – Arctic manual which covers two GFA Domains. This document forms an important part of the training program on local aviation weather knowledge for FSS working in the area and a useful tool in the day-to-day service delivery by FSS.

Within the GFA domains, the weather shows strong climatological patterns controlled either by season or topography. This manual describes the Domain of the GFACN33 (Ontario-Quebec). This area offers beautiful skies and landscapes for flying but can also provide harsh flying conditions. As most pilots flying the region can attest, these variations in weather can take place quiet abruptly. From the cold shores of Hudson Bay to the lush hills of the Eastern Townships, local topography plays a key role in determining both the general climatology and local flying conditions in a particular region.

This manual provides some insight on specific weather effects and patterns in this area. While a manual cannot replace intricate details and knowledge of Ontario-Quebec that FSS and experienced pilots of the area have acquired over the years, this manual is a collection of that knowledge taken from interviews with local pilots, dispatchers, Flight Service Specialists, and MSC personnel.

By understanding the weather and hazards in this specific area, FSS will be more able to assist pilots to plan their flights in a safe and efficient manner. While this is the manual's fundamental purpose, NAV CANADA recognizes the value of the information collected for pilots themselves. More and better information on weather in the hands of pilots will always contribute to aviation safety. For that reason, the manuals are being made available to NAV CANADA customers.



Acknowledgements

This manual was made possible through funding by NAV CANADA, Flight Information Centre project office.

NAV CANADA would like to thank The Meteorological Service of Canada (MSC), both national and regional personnel, for working with us to compile the information for each Graphic Area Forecast (GFA) domain, and present it in a userfriendly, professional format. Special thanks also go to meteorologists Ross Klock, Mountain Weather Centre, Kelowna and Gilles Simard, Environmental Weather Services Office –East, Rimouski, Quebec and John Mullock, Mountain Weather Centre, Kelowna. Ross's and Gilles's regional expertise has been instrumental for the development of the Ontario-Quebec GFA document while John's experience and efforts have ensured high quality and consistent material from Atlantic to Pacific to Arctic.

This endeavour could not have been as successful without the contributions of many people within the aviation community. We would like to thank all the participants that provided information through interviews with MSC, including flight service specialists, pilots, dispatchers, meteorologists and other aviation groups. Their willingness to share their experiences and knowledge contributed greatly to the success of this document.

Roger M. Brown January, 2002

Readers are invited to submit any comments to:

NAV CANADA Customer Service Centre 77 Metcalfe St. Ottawa, Ontario K1P 5L6

Toll free phone line: 1-800-876-4693-4 (within North America disregard the last digit)

Toll-free fax line: 1-877-663-6656

E-mail: service@navcanada.ca



TABLE OF CONTENTS

PREFACE		iii
ACKNOWLEDGE	MENTS	iv
INTRODUCTION		ix
CHAPTER 1	BASICS OF METEOROLOGY	1
	Heat Transfer and Water Vapour	1
	Lifting Processes	2
	Subsidence	3
	Temperature Structure of the Atmosphere	4
	Stability	5
	Wind	
	Air Masses and Fronts	6
CHAPTER 2	WEATHER HAZARDS TO AVIATION	9
	Introduction	9
	Icing	9
	Types of Aircraft Ice	10
	Meteorological Factors Affecting Icing	
	Aerodynamic Factors Affecting Icing	
	Other Forms of Icing	
	Visibility	
	Types of Visibility	
	Causes of Reduced Visibility	
	Wind, Shear and Turbulence	
	Stability and the Diurnal Variation in Wind	
	Wind Shear	
	Relationship Between Wind Shear and Turbulence .	22
	Low Levels Jets - Frontal	
	Low Level Jets - Nocturnal	
	Topographical Effects on Wind	
	Lee Waves	
	The Formation of Lee Waves	29
	Characteristics of Lee Waves	29
	Clouds Associated with Lee Waves	31
	Fronts	32
	Frontal Weather	33
	Frontal Waves and Occlusions	33
	Thunderstorms	35
	The Life Cycle of a Thunderstorm	36
	Types of Thunderstorms	38
	Severe Thunderstorms	
	Thunderstorm Hazards	40
	Cold Weather Operations	
	Volcanic Ash	
	Deformation Zone	45



CHAPTER 3	WEATHER PATTERNS	49
	Introduction	49
	Geography of Ontario and Quebec	49
	The Mean Atmospheric Circulation System	
	Upper Troughs and Upper Ridges	
	Semi-Permanent Surface Features	
	Migratory Systems	
	Winter Storms	
	Summer Storms	
	Alberta Lows	
	Colorado Lows	
	Great Lakes Lows	66
	Hudson Bay Lows	
	Polar Lows	
	Hatteras Lows	
	Highs	
	Tropical Depressions, Tropical Storms and Hurricanes	
	Cold Lows	
	Seasonal Migratory Birds	
CHAPTER 4	REGIONAL WEATHER & LOCAL EFFECTS	
	Introduction	73
	Ontario	74
	Local Effects	76
	Hudson Bay Coast and Lowlands Fort Severn -	
	Moosonee	76
	Fort Severn – Cape Henrietta Maria	77
	Cape Henrietta Maria – Moosonee	78
	Northwestern Ontario	80
	Red Lake – Pickle Lake - Sandy Lake –	
	Big Trout Lake	81
	Kenora – Fort Frances – Dryden -Sioux Lookout –	
	Thunder Bay	
	Northeastern Ontario	
	Kapuskasing – Timmins - Geraldton	85
	North Bay – Sudbury - Sault Ste. Marie –	
	Wawa – Marathon	
	Southern Ontario	88
	Windsor – London – Hamilton – St Catharines –	
	Buttonville – Muskoka	
	Peterborough – Trenton – Kingston – Ottawa	
	The Great Lakes	
	Lake Superior	
	Lake Huron	
	Lake Erie	
	Lake Ontario	.100



	Quebec
	Nunavik
	Kangiqsualujjuaq to Kuujjuaq
	Labrador Trough (also called Schefferville Trench)106
	Riviere aux Feuilles (Leaves River)
	Tasiujuaq Area
	Aupaluk Area
	Kangirsuk
	Area from Quaqtaq to Salluit to Ivujivik
	Akulivik and Puvirnituq
	(northern section of the east shore of Hudson Bay)
	Inukjuak
	Umiujaq
	James Bay Region
	Local effects - Coastal Area
	Dams Area
	Northwestern Quebec
	Chibougamau Mistassini Area
	Abitibi Area
	Otis Range and White Mountains Areas
	Saguenay, Lake Saint-Jean, and surrounding
	mountainous terrain
	Local effects - Saguenay River Valley
	Lake St. Jean Area
	5
	Laurentides Wildlife Reserve
	Laurentides Wildlife Reserve
	Laurentides Wildlife Reserve
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135
	Laurentides Wildlife Reserve
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.138
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area.136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139Eastern Townships and Beauce.139
	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139Eastern Townships and Beauce.139Eastern Townships Area.140
CHADTED 5	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139Eastern Townships and Beauce.139Eastern Townships Area.140Beauce Area.141
CHAPTER 5	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139Eastern Townships and Beauce.139Eastern Townships Area.140Beauce Area.141AIRPORT CLIMATOLOGY.145
GLOSSARY	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139Eastern Townships and Beauce.139Eastern Townships Area.140Beauce Area.141AIRPORT CLIMATOLOGY.145
GLOSSARY TABLE OF SYMBO	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139Eastern Townships and Beauce.139Eastern Townships Area.140Beauce Area.141AIRPORT CLIMATOLOGY.145OLS.211
GLOSSARY	Laurentides Wildlife Reserve.128Mount Valin Highlands.129Saint Lawrence river valley and surrounding terrain.131Local effects - Greater Montreal and Monteregie.132Lower Laurentians.135Quebec City - Trois-Rivieres - Drummondville Area136Greater Quebec City Area.136Drummondville.137Trois-Rivieres.137Middle Saint Lawrence Estuary Area.138South Shore Area.139Eastern Townships and Beauce.139Eastern Townships Area.140Beauce Area.141AIRPORT CLIMATOLOGY.145



viii




Introduction

Meteorology is the science of the atmosphere, a sea of air that is in a constant state of flux. Within it storms are born, grow in intensity as they sweep across sections of the globe, then dissipate. No one is immune to the day-to-day fluctuations in the weather, especially the aviator who must operate within the atmosphere.

Traditionally, weather information for the aviation community has largely been provided in textual format. One such product, the area forecast (FA), was designed to provide the forecast weather for the next twelve hours over a specific geographical area. This information consisted of a description of the expected motion of significant weather systems, the associated clouds, weather and visibility.

In April 2000, the Graphical Area Forecast (GFA) came into being, superceding the area forecast. A number of MSC Forecast Centres now work together, using graphical software packages, to produce a single national graphical depiction of the forecast weather systems and the associated weather. This single national map is then partitioned into a number of GFA Domains for use by Flight Service Specialists, flight dispatchers and pilots.



NAV CANAD

This Ontario-Quebec Local Area Knowledge Aviation Weather Manual is one of a series of six similar publications. All are produced by NAV CANADA in partnership with the MSC. These manuals are designed to provide a resource for Flight Service Specialists and pilots to help with the understanding of local aviation weather. Each of the six manuals corresponds to a Graphical Area Forecast (GFA) Domain, with the exception of the Nunavut - Arctic manual which covers two GFA Domains. MSC aviation meteorologists provide most of the broader scale information on meteorology and weather systems affecting the various domains. Experienced pilots who work in or around it on a daily basis, however, best understand the local weather. Interviews with local pilots, dispatchers and Flight Service Specialists, form the basis for the information presented in Chapter 4.

Within the domains, the weather shows strong climatological patterns that are controlled either by season or topography. For example, in British Columbia there is a distinctive difference between the moist coastal areas and the dry interior because of the mountains. The weather in the Arctic varies strongly seasonally between the frozen landscape of winter and the open water of summer. These changes are important in understanding how the weather works and each book will be laid out so as to recognize these climatological differences.

This manual describes the weather of the GFACN33 (Ontario-Quebec). This area often has beautiful flying weather but can also have some of the harshest flying conditions in the world. As most pilots flying in the region can attest, these variations in flying weather can take place quite abruptly. From the Northern Ontario to the Eastern Townships of Quebec, local topography plays a key role in determining both the general climatology and local flying conditions in a particular region. Statistically, approximately 30% of aviation accidents are weather related and up to 75% of delays are due to weather.

This manual is "instant knowledge" about how the weather behaves in this area but it is not "experience". The information presented in this manual is by no means exhaustive. The variability of local aviation weather in Ontario and Quebec could result in a publication many times the size of this one. However, by understanding some of the weather and hazards in these areas, pilots may be able to relate the hazards to topography and weather systems in areas not specifically mentioned.



Chapter 1

Basics of Meteorology

To properly understand weather, it is essential to understand some of the basic principles that drive the weather machine. There are numerous books on the market that describe these principles in great detail with varying degrees of success. This section is not intended to replace these books, but rather to serve as a review.

Heat Transfer and Water Vapour

The atmosphere is a "heat engine" that runs on one of the fundamental rules of physics: excess heat in one area (the tropics) must flow to colder areas (the poles). There are a number of different methods of heat transfer but a particularly efficient method is through the use of water.

Within our atmosphere, water can exist in three states depending on its energy level. Changes from one state to another are called phase changes and are readily accomplished at ordinary atmospheric pressures and temperatures. The heat taken in or released during a phase change is called latent heat.




Fig. 1-1 - Heat transfer and water vapour

How much water the air contains in the form of water vapour is directly related to its temperature. The warmer the air, the more water vapour it can contain. Air that contains its maximum amount of vapour, at that given temperature, is said to be saturated. A quick measure of the moisture content of the atmosphere can be made by



looking at the dew point temperature. The higher (warmer) the dew point temperature, the greater the amount of water vapour.

The planetary heat engine consists of water being evaporated by the sun into water vapour at the equator (storing heat) and transporting it towards the poles on the winds where it is condensed back into a solid or liquid state (releasing heat). Most of what we refer to as "weather," such as wind, cloud, fog and precipitation is related to this conversion activity. The severity of the weather is often a measure of how much latent heat is released during these activities.

Lifting Processes

The simplest and most common way water vapour is converted back to a liquid or solid state is by lifting. When air is lifted, it cools until it becomes saturated. Any additional lift will result in further cooling which reduces the amount of water vapour the air can hold. The excess water vapour is condensed out in the form of cloud droplets or ice crystals which then can go on to form precipitation. There are several methods of lifting an air mass. The most common are convection, orographic lift (upslope flow), frontal lift, and convergence into an area of low pressure.




Fig. 1-2 - Convection as a result of daytime heating




Fig.1-3 - Orographic (upslope) lift






Fig.1-4 - Warm air overrunning cold air along a warm front




Fig. 1-5 - Divergence and convergence at the surface and aloft in a high low couplet

Subsidence

Subsidence, in meteorology, refers to the downward motion of air. This subsiding motion occurs within an area of high pressure, as well as on the downward side of a range of hills or mountains. As the air descends, it is subjected to increasing atmospheric pressure and, therefore, begins to compress. This compression causes the air's temperature to increase which will consequently lower its relative humidity. As a result, areas in which subsidence occurs will not only receive less precipitation than surrounding areas (referred to as a "rain shadow") but will often see the cloud layers thin and break up.






Fig.1-6 - Moist air moving over mountains where it loses its moisture and sinks into a dry subsidence area

Temperature Structure of the Atmosphere

The temperature lapse rate of the atmosphere refers to the change of temperature with a change in height. In the standard case, temperature decreases with height through the troposphere to the tropopause and then becomes relatively constant in the stratosphere.

Two other conditions are possible: an inversion, in which the temperature increases with height, or an isothermal layer, in which the temperature remains constant with height.




Fig. 1-7 - Different lapse rates of the atmosphere





The temperature lapse rate of the atmosphere is a direct measurement of the stability of the atmosphere.

Stability

It would be impossible to examine weather without taking into account the stability of the air. Stability refers to the ability of a parcel of air to resist vertical motion. If a parcel of air is displaced upwards and then released it is said to be unstable if it continues to ascend (since the parcel is warmer than the surrounding air), stable if it returns to the level from which it originated (since the parcel is cooler than the surrounding air), and neutral if the parcel remains at the level it was released (since the parcel's temperature is that of the surrounding air).

The type of cloud and precipitation produced varies with stability. Unstable air, when lifted, has a tendency to develop convective clouds and showery precipitation. Stable air is inclined to produce deep layer cloud and widespread steady precipitation. Neutral air will produce stable type weather which will change to unstable type weather if the lifting continues.





The stability of an air mass has the ability to be changed. One way to destabilize the air is to heat it from below, in much the same manner as you would heat water in a kettle. In the natural environment this can be accomplished when the sun heats the ground which, in turn, heats the air in contact with it, or when cold air moves over a warmer surface such as open water in the fall or winter. The reverse case, cooling the air from below, will stabilize the air. Both processes occur readily.

Consider a typical summer day where the air is destabilized by the sun, resulting in the development of large convective cloud and accompanying showers or thundershowers during the afternoon and evening. After sunset, the surface cools and the air mass stabilizes slowly, causing the convective activity to die off and the clouds to dissipate.



On any given day there may be several processes acting simultaneously that can either destabilize or stabilize the air mass. To further complicate the issue, these competing effects can occur over areas as large as an entire GFA domain to as small as a football field. To determine which one will dominate remains in the realm of a meteorologist and is beyond the scope of this manual.

Wind

Horizontal differences in temperature result in horizontal differences in pressure. It is these horizontal changes in pressure that cause the wind to blow as the atmosphere attempts to equalize pressure by moving air from an area of high pressure to an area of low pressure. The larger the pressure difference, the stronger the wind and, as a result, the day-to-day wind can range from the gentlest breeze around an inland airfield to storm force winds over the water.




Fig. 1-9 - The greater pressure changes with horizontal difference, the stronger the wind

Wind has both speed and direction, so for aviation purposes several conventions have been adopted. Wind direction is always reported as the direction from which the wind is blowing while wind speed is the average steady state value over a certain length of time. Short-term variations in speed are reported as either gusts or squalls depending on how long they last.

Above the surface, the wind tends to be relatively smooth and changes direction and speed only in response to changes in pressure. At the surface, however, the wind is affected by friction and topography. Friction has a tendency to slow the wind over rough surfaces whereas topography, most commonly, induces localized changes in direction and speed.

Air Masses and Fronts

Air Masses

When a section of the troposphere, hundreds of miles across, remains stationary or moves slowly across an area having fairly uniform temperature and moisture, then the



air takes on the characteristics of this surface and becomes known as an air mass. The area where air masses are created are called "source regions" and are either ice or snow covered polar regions, cold northern oceans, tropical oceans or large desert areas.

Although the moisture and temperature characteristics of an air mass are relatively uniform, the horizontal weather may vary due to different processes acting on it. It is quite possible for one area to be reporting clear skies while another area is reporting widespread thunderstorms.

Fronts

When air masses move out of their source regions they come into contact with other air masses. The transition zone between two different air masses is referred to as a frontal zone, or front. Across this transition zone temperature, moisture content, pressure, and wind can change rapidly over a short distance.

The principal types of fronts are:

Cold Front - The cold air is advancing and undercutting the warm air. The leading edge of the cold air is the cold front.	A A	A A
Warm front - The cold air is retreating and being replaced by warm air. The trailing edge of the cold air is the warm front.	عمد	"The
Stationary front - The cold air is neither advancing nor retreating. These fronts are frequently referred to quasi-stationary fronts although there usually is some small-scale localized motion occurring.	×	x
Trowal - Trough of warm air aloft.	1	1

Table 1-1

